


Vascular Surgery


Open Abdominal Surgery

Open abdominal surgery involves making a small incision to repair an aortic aneurysm or blockage of the aorta. In many cases, A cylinder like tube called a vascular graft may be used to repair the aneurysm or stenosis.




Angioplasty and Stenting

Angioplasty and stenting are a minimally invasive procedure in which your surgeon opens a narrowed artery using a catheter-guided balloon or vascular stent. This method can be used to treat patients with peripheral artery disease and allows vascular access for patients who need dialysis. This procedure treats conditions that include peripheral artery disease, a narrowing of the blood vessels that carry oxygenated blood to the arms and leg.



Thrombectomy and Endarterectomy


A thrombectomy is a procedure to remove a blood clot from an artery. It is done to restore proper blood flow and prevent tissue necrosis. Angioplasty and stenting may also be performed at the same time as a thrombectomy. Endarterectomy involve the surgical removal of plaque within the inner lining of the arteries feeding blood to your arms or legs, respectively.



Vascular Surgery


Vascular Bypass Surgery

Bypass grafting is a surgical treatment that bypass the stenosis or occlusion area using artificial graft or vein graft to maintain blood flow. This surgery may be used to treat patients with the following disorders: Peripheral artery disease, affecting blood flow to the arms and legs.



Arteriovenous (AV) Fistula

During an AV fistula procedure, your vascular surgeon connects a vein in your forearm directly to an artery. This makes the vein stronger and wider, and more easily to be accessed for patients with kidney failure who require dialysis. When cephalic veins are very thin, artificial Vascular graft may be used.



Varicose Veins Surgery:

Recommendations for the relief to varicose vein symptoms include wearing support stocking and elevating your legs. If that does not support enough, a surgical procedure can be done to remove the varicose veins or seal off under local anesthesia. Liquid preparations are used for the treatment of small nonaxial veins with cosmetic concerns.

